This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

IMMUNO-ANALYTICAL METHOD FOR AMPHETAMINES AND REAGENTTHEREOF

Patent Number:

JP56125666

Publication date:

1981-10-02

Inventor(s):

SAITO SUMIYASU; others: 07

Applicant(s):

FUJISAWA PHARMACEUT CO LTD

Requested Patent

☐ JP56125666

Application Number: JP19800029460 19800307

Priority Number(s):

IPC Classification:

G01N33/54

EC Classification:

Equivalents:

Abstract

PURPOSE:To discriminately detect the amphetamine and methamphetamine in humor by the antibody obtained by sensitizing the high polymer material bound on the benzene ring of amphetamines with warmblooded animal.

CONSTITUTION: The composite of a high polymer material, for example, (P)-N= N-(Q) or (P)-(CH2)nNH-(Q) bound on the benzene ring through a spaur for providing immunogenic to the amphetamines expressed by the formula (I) and a high polymer material is injected subcutaneously or muscularly into warm-blooded animal, and after a fixed period, the blood of said animal is sampled, whereby the antibody is obtained. The immuno-analytical method of the amphetamines obtained in this way permits straight application of known methods, and has virtually no cross reactivity with the methamphetamine of the amphetamine antibody.